



**ORIGINAL ARTICLE**

**Limnological Assessment of Keetham Lake at Soor-Sarovar Bird Sanctuary in Agra**

**Vishwakant**

Department of Zoology, Agra College, Agra

Email: [gupta.vishwakant2@gmail.com](mailto:gupta.vishwakant2@gmail.com)

**ABSTRACT**

*To investigate the limnology of fresh water Keetham Lake in Taj city of Agra (U.P.), samples of water collected from 3 different stations at each month from March 2013 to February 2014, the eleven (11) physical and chemical factors taken for the study of Keetham Lake. These limnological parameters were estimated and analyzed systematically. Water temperature, Turbidity, Light penetration, Depth, pH values varied from 16 to 35° C, 72 to 142 cm, 20 to 28.8 cm., 12 to 22.5 ft, 6.8 to 8.5 respectively. A value of D.O. was 5.5 to 8.4 mg/l, B.O.D. was 63-139.3 mg/l and 85-220 mg/l, C.O.D. was recorded. CO<sub>2</sub> and alkalinity were limited from 27.0 to 39.5 mg/l and 185.87 to 315.0 mg/l respectively. T.D.S. was recorded in bearable concentrations in such Lake. Lake never revealed any significant pollution at hypothetical stations as well as no significant variations was observed among various parameters under study. Correlation coefficients were found out to make correlation analysis among parameters. At a glance present Research work showed that all stations of Lake had excellent level of D.O., alkaline pH, moderate alkalinity, enriched nutrients, better transparency which supported abundance of biotic flora and fauna. The Lake was also found to have excellent place for migratory birds.*

**Key words:** limnology, pollutometric, Keetham, correlation analysis

*Received: 5<sup>th</sup> September, 2016, Revised: 15<sup>th</sup> October 2016, Accepted: 22<sup>nd</sup> October 2016*

*©2016 Council of Research & Sustainable Development, India*

**How to cite this article:**

Vishwakant (2016): Limnological Assessment of Keetham Lake at Soor-Sarovar Bird Sanctuary in Agra. *Annals of Natural Sciences*, Vol. 2[4]: December, 2016: 22-27.

**INTRODUCTION**

In India greater regional differences are found in the topography, ecology, climate etc. As a result of these differences the limnological resources (Lake, pond & reservoirs) and their morphometries are very unevenly distributed. Nevertheless India is blessed with vast water resources in the form of rivers, estuaries, Lakes, natural and artificial ponds and mangrove wetlands, which are ideal natural habitat for biotic placements and sites for fishery and its development.

Physico-chemical state of any water body depends on natural contents of physical factors & nutrients. Physico-chemical equilibrium is must between components of water body dissolved in it for fishery development and conservation of biodiversity. Therefore any seasonal, periodic (natural) and anthropogenic, environmental (artificial) alteration in water characteristics like temperature, light, wind current, turbidity, D.O., pH, alkalinity, total hardness, other chemicals nutrients and soil texture & composition, diversity and food chain of aquatic ecosystems and even deterioration of water resources if physico-chemical features get distributed more.. This is an exercise in identifying limnological conditions of Keetham Lake, although Lake is considered a best asylum for birds from abroad. They come to Keetham Lake from a long distance during their longer travel as migration. However author has not taken this account for his research in present research piece of work. The proposed work has, therefore, been undertaken, keeping in view the complete survey with the aim of knowing abiotic parameters of such Lake.

Present limnological investigations of Keetham Lake are undertaken to record monthly fluctuations in abiotic parameters (temperature, light penetration, depth, turbidity, hydrogen ion concentration, dissolved oxygen, free carbon dioxide, alkalinity, BOD, COD etc), for a period of one year only. However the vital processes are at the optimum when the physical and chemical conditions are normal. The eleven (11) physical and chemical factors have taken for the study of Keetham Lake.

The important physical and chemical factors taken for the study of Keetham Lake water play a significant role in controlling the distribution and growth of biotic flora and fauna, suitability for migratory birds, use the water for other purposes including recreational activities.

#### MATERIAL AND METHODS

Water samples were collected help of labor in the second week of each month from March 2013 to February 2014. Three replicates each of 2 lit. Samples were collected in sampling bottles in the morning hours from three 'I', 'II' & 'III' sampling stations and brought to the laboratory for analysis. Standard methods for the examination of water and waste water (APHA 1996) were used for the analysis.

At the time of water sampling (at spots) temperature, depth and light penetration were recorded. Temperature was recorded with the help of mercury Thermometer. For the measurement of D.O., water samples were fixed with the help of a manganous sulphate and alkali iodideazide solution at the sites and analyzed in the laboratory using Winkler's iodo metric method. Turbidity was measured by digital nephelometer, B.O.D. and C.O.D. were analyzed by B.O.D. incubator and potassium- dichromate methods respectively. pH (using a pH meter), CO<sub>2</sub> (using Sodium carbonate titration), alkalinity (titration method by H<sub>2</sub>SO<sub>4</sub> and NaOH using phenolphthalein and methyl orange indicator), chloride (by Mohr's argentometric method using potassium chromate as indicator), Ammonia (by simple nesslerisation method), phosphate (by Atkins modification, stannous chloride methods), Silicate (Ammonium molybdate method), T.D.S. (by evaporation method), and nitrate using phenol-disulphonic acid colorimetric test.

**Table 1:** Limnological parameters with their mean values and Standard Deviations of Keetham Lake (mean values shown in the table taken from 3 stations as I, II and III) (March 2013 to Feb. 2014)

Parameter/ month	Water temp.	Turbidity	Light penetration	Depth (feet)	pH	D.O.
March	27.0 ± 0.69	85 ± 1.06	28.8 ± 0.98	14 ± 0.77	7.8 ± 0.33	7.7 ± 0.78
April	30.4 ± 0.91	72 ± 1.40	26.8 ± 0.90	14.5 ± 0.50	8.0 ± 0.16	6.2 ± 0.88
May	34.6 ± 0.66	74 ± 1.70	24.0 ± 0.95	12.0 ± 0.28	8.4 ± 0.48	5.5 ± 0.26
June	32.8 ± 0.57	85 ± 1.0	24.1 ± 1.39	13.5 ± 0.40	8.5 ± 0.32	6.2 ± 0.18
July	33.0 ± 1.21	118 ± 0.81	21.5 ± 0.84	19.0 ± 0.47	6.8 ± 0.28	6.2 ± 0.34
Aug.	30.6 ± 0.46	125 ± 0.93	20.7 ± 0.54	22.5 ± 0.64	7.5 ± 0.38	6.8 ± 0.18
Sep.	29.2 ± 0.89	142 ± 0.48	20.0 ± 0.93	22.0 ± 0.52	7.9 ± 0.87	7.3 ± 0.86
Oct.	26.0 ± 1.44	125 ± 0.85	22.5 ± 1.02	20.0 ± 0.48	8.0 ± 0.28	7.8 ± 0.65
Nov.	23.4 ± 0.86	98 ± 1.04	25.0 ± 1.00	20.2 ± 0.35	8.1 ± 0.76	7.1 ± 0.34
Dec.	18.2 ± 0.75	98 ± 1.30	28.2 ± 0.94	18.5 ± 0.76	8.0 ± 0.56	8.2 ± 0.76
Jan.	16.7 ± 0.88	88 ± 0.82	26.0 ± 0.90	19.2 ± 0.52	7.9 ± 0.45	8.1 ± 0.64
Feb.	19.6 ± 0.94	78 ± 0.85	27.8 ± 0.9	18.5 ± 0.88	7.7 ± 0.58	8.4 ± 0.36

All Mean values derived from three stations viz. I, II & III are expressed in mg/lit., except Temp. (°C), pH, Turbidity and Light penetration (cm), Depth (feet). All values are given with their Standard Deviations (S.D.) respectively. Higher S.D. shows that values have more difference in three stations while Lower S.D. shows that values have less difference among three stations.

**Table 2:** Limnological parameters with their mean values and Standard Deviations of Keetham Lake (mean values shown in the table taken from 3 stations as I, II and III) (March 2013 to Feb. 2014)

Parameter/ month	B.O.D.	C.O.D.	CO <sub>2</sub>	Alkalinity	TDS
March	88 ± 6.6	130 ± 2.10	34.0 ± 0.67	255.0 ± 4.3	575 ± 3.8
April	97.6 ± 6.12	145 ± 3.87	30.0 ± 0.86	275.0 ± 3.6	854 ± 3.5
May	102.3 ± 7.13	170 ± 2.1	27.0 ± 0.65	280.2 ± 3.2	1090 ± 2.7
June	116.5 ± 6.2	185 ± 2.90	29.0 ± 0.34	315.0 ± 3.1	1640 ± 7.5
July	121.3 ± 4.7	192 ± 5.35	39.5 ± 0.98	185.7 ± 5.3	1820 ± 7.7
Aug.	139.3 ± 5.7	220 ± 4.67	36.5 ± 0.88	194.5 ± 2.8	1520 ± 4.5
Sep.	96.2 ± 4.9	210 ± 3.98	30.8 ± 1.1	208.5 ± 3.6	1010 ± 4.8
Oct.	98 ± 8.1	188 ± 4.55	28.5 ± 1.3	214 ± 3.7	1100 ± 7.2
Nov.	84 ± 4.8	152 ± 5.6	29.8 ± 0.46	225.0 ± 4.8	1250 ± 6.8
Dec.	65.6 ± 4.9	85 ± 5.80	31.6 ± 0.55	238.2 ± 2.8	1140 ± 6.3
Jan.	63.0 ± 5.09	94 ± 3.88	28.0 ± 0.90	245.0 ± 2.6	1210 ± 5.6
Feb.	77.6 ± 4.4	89 ± 4.1	30.0 ± 0.94	250.0 ± 3.1	900 ± 4.8

### CORRELATION ANALYSIS

A positive and significant correlation was found out between water temperature, BOD, COD, CO<sub>2</sub>, and Chloride. A negative and significant correlation has been recorded between temperature and D.O. Turbidity exhibited negative relation with light penetration, pH and D.O. while positive correlation with most of parameters. pH showed positive correlation with alkalinity, chloride and D.O. and negative correlation with rest of the parameters. D.O. positively correlated with light penetration and pH. BOD and COD had negative correlation with light penetration, D.O., and chloride while other parameter more or less positively correlated with both ones. CO<sub>2</sub> negatively significant correlated with D.O. and alkalinity in short strength, while positively correlated with most of the parameters.

### RESULTS AND DISCUSSION

#### WATER TEMPERATURE:

The water temperature governs species richness and diversity (Thirumalaet *al.*, 2006) distribution and their rates of activity besides water mixing, turbulence and production of currents (Ruttner, 1963; Cole, 1983). Maximum 34.6°C recorded in May 2013, while minimum 16.7°C in Jan. 2014. Variation in water temperature in present work as monthly changing was found due to heat gain from solar radiation from the air and transfer by conduction in water body. At different water level temperature was observed uniform, so no thermal stratification was noted, which made Lake productive. A positive correlation between bright sunlight, its longevity and water temperature was observed by Ahmad & Singh, 1990; Vishwakant, 2007.

#### TURBIDITY:

Turbidity makes the water bodies unfit for direct or indirect consumption by human for various uses. Excess value of turbidity hinders the fish yield by reducing other factors like light penetration, pH, D.O., ammonia, chlorides etc. (Vishwakant, 2007). At station 'I' maximum 146 recorded in September 2013, while minimum 76 in April 2013 at station 'III'. However mean values ranged from 72 to 142 cm. Turbidity is the Pollute metric factor. It was observed in high proportion at station 'I' (146 cm in Sept 2013) because Domestic discharge, dirt other impurities coming from nearby area also anthropogenic and zoogenic activities, inflow of silt, surface run off during raining season raised turbidity. In present work higher turbidity during monsoon due to suspended particles might observe nutrient concentration (phosphate & nitrate) making them unavailable for plankton production. While minimum turbidity was noted in summer. Turbidity was found as inversely related with light penetration, pH, and D.O.

**LIGHT PENETRATION (TRANSPARENCY):**

Maximum average 28.8 recorded in March 2013, while minimum 20.0 in Sept. 2013. Maximum transparency was recorded in spring season; moderate in winter as well. High transparency could be attributed to less decomposition of organic matter due to less temperature and less input of solids & wastes. Low transparency and high turbidity reduce photosynthesis of plankton & plants which in turn lessen the growth of biota & density of fish.

**DEPTH:**

At station 'II' maximum 24.5 ft recorded in Aug. 2013, while minimum 13.5 ft ( station III) in May 2013. Average values were from 12 to 22.5 ft. Although depth was maximum observed in rainy season, but higher values of depth were also noted in winter & spring. An inverse relation was observed between Depth and light penetration; In present work all stations had saturated water level for suitability for Lake to exist.

**pH:**

The pH an indicative of hydrogen ion concentration, the hydrogen ion concentration of natural waters is very essential environmental factor, the variations of which, among other causes are linked with species composition and life processes of animal and plants communities' inhabiting them (Jhingran, 1988). PH is often used as an index for water condition. Mean values from minimum  $6.8 \pm 0.28$  to maximum  $8.5 \pm 0.32$  recorded in July and June 2013 respectively. The lower values of pH during rainy season in present study got support by the works of Kaushik, *et al.*, 1991; Saxena 1994. Moderate value of pH in water spring and reached max. in summer might be due to higher photosynthesis, algal blooms, reach population of submerged macrophytes as reported by Saxena 1994.

In all stations in present work pH values did not undergo any sharp variation during whole of the year. Low records of pH during raining season was observed due to lower transparency, higher turbidity, and more depth, higher CO<sub>2</sub> lower D.O., reduced photosynthesis, which ultimately reduced fish production in such season. Moderate value of pH was observed in winter- spring seasons which led to flourishing of bio diversity and fruitful settlements of migratory birds.

**DISSOLVED OXYGEN (D.O.):**

Mean values from minimum  $5.5 \pm 0.26$  to maximum  $8.4 \pm 0.36$  recorded in May 2013 and Feb. 2014 respectively. D.O. directly affects survival, development, density and distribution of biota in an ecosystem. (Vijayakumar, *et al.*, 1999). In fact, at any time the concentration of D.O. is the balance between oxygen produced by photosynthesis during day and oxygen lost to respiring flora and fauna day and night. Higher values were recorded in winter and spring seasons due to less microbial decomposition of dead organic matter, its high solubility at low temperature (Kant and Raina, 1990; Sulthara, 2005), holding capacity of water, moderate photosynthesis and raised growth of submerged macrophytes, (Vishwakant, *et al.*, 2007) which led to good fish yield. While lower values of D.O. were observed in summer & rainy season due to higher free CO<sub>2</sub>, lower photosynthesis, high Turbidity, higher respiration of biota, higher microbial activity, organic pollutants. All the time D.O. was observed in significant concentrations which always helped the Lake to keep promoting life conditions for existing biological organisms and visitor migratory birds.

**B.O.D.:**

Mean values from minimum  $63.0 \pm 5.09$  to maximum  $139.3 \pm 5.7$  recorded in Jan. 2014 and Aug. 2013 respectively.

**C.O.D.:**

Mean values from minimum  $85.0 \pm 5.8$  to maximum  $220 \pm 4.6$  recorded in Dec. and Aug. 2013 respectively.

**CO<sub>2</sub>:**

The presence and absence of CO<sub>2</sub> is determined by its consumption by algae and plants during photosynthesis and also through its diffusion from air. Mean values from minimum  $27.0 \pm 0.65$  to maximum  $39.5 \pm 0.98$  recorded in May and July 2013 respectively. It was found that CO<sub>2</sub> inversely correlated with D.O. in different seasons. Higher values in summer rainy season were observed due to higher decomposition of dead organic matter & submerged macrophytes, lower photosynthesis, cloudy weather and increased suspended matter etc. led to lower biological activities, while in winter spring season lower CO<sub>2</sub> were observed in all stations, which led to higher growth of biotic communities and excellent water quality.

**ALKALINITY:**

The label of alkalinity is a measure of productivity of water. (Ghosh and George, 1989; Sharma and Kaushal, 2004). Mean values from minimum  $185.7 \pm 5.3$  to maximum  $315 \pm 3.1$  recorded in July and June 2013 respectively.

Alkalinity in natural water is caused by salts of carbonates, bicarbonate, hydroxide etc. of course carbonate and bi carbonate forms and index of total alkalinity have a direct wearing and liner relationship with pH (Vishwakant, 2010). An increased pH in Keetham Lake due to carbonates corroborates with the work of (Hutchinson 1957; Jhingran 1988; Goldman and Horne 1983). In all stations alkalinity had excellent values resulting their productive nature especially coupled with high production of phytoplankton (although known plankton analysis involved in given parameters). This is important factor to promote life conditions for existing biological organisms and visitor migratory birds.

**CONCLUSION**

In the present paper all stations of Lake Keetham limnological parameters have been found to have changed during study. Present work emphasized over changed values of parameters during different months of investigated period, due to sewage, organic, agriculture, sedimental, siltation pollutants which ultimately changed the biotic profile of communities. All the parameters studied were found in the optimum range to sustainable development of this Lake. All the parameters from water temperature to TDS under investigation were in significant concentrations giving excellent water quality to the Lake which finally promoted various feasible life conditions for existing biological organisms and visitor migratory birds. A proper & systematic development of Lake could lead to fisheries and allied sectors. The water of Lake can be used for domestic (washing), agriculture, aquaculture and recreational purposes. So author selected such larger Lake for assessment of its hydro biological characteristics.

**REFERENCES**

1. Ahmed S.H. and Singh 1990. Limnological features of Kawar Lake Bihar. Seminar on Wetland Ecology and management at Keoladeo nat. park, Bharatpur 23-25 Feb. 1990.
2. APHA (1996): Standard methods for the examination of water and waste water; 18<sup>th</sup> ed., APHA, AWWA, WPCF, New York.
3. Cole G.A. (1983): Text book of limnology. (3<sup>rd</sup> ed.) C.V. Mosby Company, London. pp: 401.
4. Ghosh A. and George J.P. (1989): Studies on the abiotic factors and zooplankton in a polluted urban reservoir, HussainSagar, Hyderabad: Impact on water quality and embryonic development of fishes. Indian J. Environ. Hlth., 37: 49-59.
5. Goldman C.R. and Horne A.J. (1983): limnology. Mc-Graw Hill International Book Company, London, pp: 464.
6. Hutchinson G.E. (1957): A treatise on limnology. Vol.1 Johnwiley and sons, New York, pp: 1015.
7. Jhingran (1988): Fish and fisheries of India. Hindustan publishing corporation (India) Delhi. pp: 954.
8. Kant S. and Raina A.K. (1990): Limnological studies of two ponds in Jammu. II, physico-chemical parameters. J. Environ. Biol., 11(2): 137-144.
9. Kaushik S., Agarkar M.S. and Saksena D.N. (1991): On the planktonic algae of sewage fed Vinek Nagar Pond at Gwalior. Poll. Res., 10(1): 25-32.
10. Ruttner R.F. (1963): Fundamentals of limnology. University of Toronto Press.

11. Saxena R.S. (1994): Ph.D. thesis, Agra University, Agra, pp: 191.
12. Sharma V.K. and Kaushal D.K. (2004): Ecological based Management of Urmila Sagar reservoir of Rajasthan. *App. Fish. & Aqua.* 4(2): 47-50.
13. Sulthara S.S. 2005. Physico-chemical properties of some fresh water bodies of Hanumangarh and Sriganganagar Districts of Rajsthan. *Poll. Res.*, 24(3): 695-698.
14. Thiruumala S., Kiran B.R., Puttaiah E.T., Kumara V. and Shakher T.R.S. (2006): Hydro chemical characteristics of Ayyanakere Lake. Near western ghats of Chikmanglore, Karnataka, *J. Aqua. Biol.*, 21(1): 111-117.
15. Vijayakumar K., Holkar D. and Kaur K. (1999): Limnological studies on Chandrampalli reservoir, Gulbaranga, In: *Freshwater Ecosystems of India.* (Ed. K. vijayakumar) Daya publishing house, Delhi, pp: 59-108.
16. Vishwakant (2007): Study of the limnological properties of some fresh water bodies of Etah district (U.P.) with reference to the effect of certain pollutants on fish productivity. Ph.D. thesis, Dr. B.R.A. university, Agra, pp: 236.
17. Vishwakant (2010): Study of nutrient status in rural pond Shikehara, Ganjdundwara (District K.R. Nagar) U.P. Rural development of India, center for research studies, Gayatry Publication, Rewa (M.P.), pp 291-298.
18. Vishwakant, Verma R.C. and Saxsena R.S. (2007): Study of some limnological properties of Harchandpur pond, District Etah (U.P.) India. *Curr. Worl. Environ.*, 2(1): 113-115.